



## Prevalence of *Capillaria plica* in Danish wild carnivores

Petersen, Heidi Huus; Nielsen, Stine Thorsø; Larsen, Gitte; Holm, Elisabeth; Chriél, Mariann

*Published in:*  
International Journal for Parasitology: Parasites and Wildlife

*Link to article, DOI:*  
[10.1016/j.ijppaw.2018.09.006](https://doi.org/10.1016/j.ijppaw.2018.09.006)

*Publication date:*  
2018

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Petersen, H. H., Nielsen, S. T., Larsen, G., Holm, E., & Chriél, M. (2018). Prevalence of *Capillaria plica* in Danish wild carnivores. *International Journal for Parasitology: Parasites and Wildlife*, 7(3), 360-363.  
<https://doi.org/10.1016/j.ijppaw.2018.09.006>


---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# Prevalence of *Capillaria plica* in Danish wild carnivores

Heidi H. Petersen\*, Stine T. Nielsen, Gitte Larsen, Elisabeth Holm, Mariann Chriél

Section for Diagnostics and Scientific Advice, National Veterinary Institute, Technical University of Denmark, Kemitorvet, 2800, Kgs. Lyngby, Denmark

## ARTICLE INFO

### Keywords:

*Capillaria plica*  
Bladderworm  
Wild carnivores  
Reservoir hosts

## ABSTRACT

*Capillaria plica* is a parasitic nematode belonging to the family Capillariidae. The adult parasites reside in the urinary tract of wild and domestic canines. The infection is most often asymptomatic, but can cause a wide range of symptoms including urinary bladder inflammation, pollakisuria, dysuria and hematuria. Canines acquire the infection by ingesting the intermediate host, the earthworm (Lumbricidae). Epidemiological studies on *C. plica* infection in wildlife are few and only one previous Danish study examined the prevalence in red foxes, while studies on prevalence in other animals are limited. We examined the urine sediment or urinary bladder from 375 Raccoon dogs (*Nyctereutes procyonoides*), 247 red foxes (*Vulpes vulpes*), 20 beech martens (*Martes foina*), 16 wild mink (*Neovison vison*), 14 otters (*Lutra lutra*), nine European polecats (*Mustela putorius*), three European badgers (*Meles meles*) and one golden jackal (*Canis aureus*) received as a part of Danish wildlife surveillance. *Capillaria plica* was detected in 73.7% of red foxes, 20.0% of beech martens, 0.5% of raccoon dogs, and in the Golden jackal. Red foxes originating from all 5 regions of Denmark were infected, although with a significantly higher prevalence in the three regions in Jutland compared to Region Zealand.

## 1. Introduction

The bladderworm *Capillaria plica* (syn. *Pearsonoma plica*) is a threadlike nematode belonging to Capillariidae family (Butterworth and Beverley-Burton, 1980). The life-cycle is indirect, involving wild and domestic canines as final host, and earthworms (Lumbricidae) as intermediate host. The final host become infected when ingesting earthworms containing first-stage larvae (L1). The L1 develop to third-stage larvae (L3) in the small intestine, migrate to the urinary bladder and moult to adult worms within two months. Adults are 13–60 mm long and embedded in the bladder mucosa. Occasionally, adults reside in urethra and renal pelvis. The female worm lays 55–67 × 26–29 µm barrel-shaped eggs with buttons on both poles (Fig. 1). The eggs are spread to the environment with urine.

*Capillaria plica* infection is generally asymptomatic, but severe urinary bladder inflammation, glomerular amyloidosis, oedema, hyperplasia of the mucosal membranes leading to pollakisuria, dysuria and hematuria are documented in dogs (Callegari et al., 2010; Lamrta and Main, 1964; Senior et al., 1980).

In Europe, *C. plica* infection is common in red foxes with reported prevalences of 78% in Germany (Bork-Mimm and Rinder, 2011), 53% in Norway (Davidson et al., 2006) and 52% in Hungary (Sréter et al., 2003). In Denmark, the prevalence has previously been documented to 80.5% in 748 red foxes collected over a five-year period in 1997–2002

(Saeed et al., 2006). Studies on *C. plica* infections in other carnivores are scarce, and no other Danish wild carnivores have previously been examined for *C. plica*.

The purpose of this study was to carry out a nation-wide cross-sectional study of *C. plica* prevalence in Danish wild carnivores, to determine their role as reservoir host for infection in Danish dogs.

## 2. Material and methods

The urinary bladder was obtained from 375 raccoon dogs (*Nyctereutes procyonoides*), 247 red foxes (*Vulpes vulpes*), 20 beech martens (*Martes foina*), 16 mink (*Neovison vison*), 14 otters (*Lutra lutra*), nine European polecats (*Mustela putorius*), three European badgers (*Meles meles*) and one golden jackal (*Canis aureus*).

The carcasses of the animals were submitted to the National Veterinary Institute, Technical University of Denmark (DTU-VET) as a part of general wildlife surveillance in 2017. The animals were either found dead, euthanized for animal welfare reasons, or hunted. The dead carnivores were transported in sealed plastic bags at –20 °C to DTU-VET and left at –80 °C for ≥2 days to inactivate potential infective parasites before necropsy. The region was listed along with information on date of death if recorded. Urinary bladders were recovered and placed in liquid-tight plastic boxes. The bladders were carefully opened and 10 ml of urine collected. If urine was absent, the bladder was cut

\* Corresponding author.

E-mail address: [hpet@vet.dtu.dk](mailto:hpet@vet.dtu.dk) (H.H. Petersen).

<https://doi.org/10.1016/j.ijppaw.2018.09.006>

Received 29 August 2018; Received in revised form 17 September 2018; Accepted 19 September 2018

2213-2244/ © 2018 The Authors. Published by Elsevier Ltd on behalf of Australian Society for Parasitology. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).



**Fig. 1.** A typical barrel-shaped *Capillaria plica* egg in urine sediment from a red fox. The egg show a slightly pitted shell and two opercules with polar plugs. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

open ( $n = 586$ ), washed with 10 ml milliQ water and the water collected in 15 ml plastic tubes. All samples were centrifuged ( $178 \times g$ ) for 10 min, the supernatant discarded to 1.5 ml, flotation fluid added to 3 ml, and samples screened for *C. plica* eggs (Fig. 1). The egg quantity per ml was recorded.

The prevalence of *C. plica* infection per animal species, and per region for red foxes was calculated together with the 95% confidence intervals. Difference in *C. plica* prevalence for red foxes between the five regions was determined by a binary logistic regression model with *C. plica* infection as the dependent variable and region of origin as the independent variable. A  $p$ -value of 0.05 was considered significant.

### 3. Results

*Capillaria plica* eggs were isolated from red foxes, raccoon dogs, beech martens and the golden jackal. Mink, otters, European polecats and European badgers were negative for *C. plica*. In total, 182 red foxes harboured *C. plica*, corresponding to a prevalence of 73.7%. Of the 375 examined raccoon dogs, only two were positive, corresponding to a prevalence of 0.5%. Of the 20 beech martens examined, four were positive (20%).

Table 1 shows the egg load (eggs per ml) in the urine or urinary bladder of positive animals ( $n = 189$ ). More than half of the positive red foxes ( $n = 94$ , 52.8%) had an egg load above 30 eggs per ml.

Fig. 2 show the prevalence of *C. plica* positive red foxes in the five Danish regions. In the three regions of Jutland (north Denmark region, central Denmark region and region of southern Denmark), *C. plica* prevalence was above 75%, while the prevalence in red foxes from the two regions on Zealand (region Zealand and capital region of Denmark) it was below 50% (Fig. 2). However, only in region Zealand, the *C. plica*

prevalence was significant lower ( $p < 0.05$ ) than the prevalence in the three regions in Jutland.

### 4. Discussion

In this cross-sectional study of *C. plica* infection in Danish wild carnivores, we identified red foxes as the most likely reservoir host for infection in dogs, as *C. plica* infection was common in red foxes (73.7%). In contrast, *C. plica* infection was rare in raccoon dogs, and absent in wild mink, otters, polecats and badgers. Lastly, we report *C. plica* infection in beech martens for the first time.

The high prevalence in red foxes in our study corresponds with the previous finding in Denmark (80.5%), although data were collected 16–21 years ago (Saeed et al., 2006). This demonstrates that *C. plica* continues to be highly prevalent in Danish red foxes and that red foxes most likely are the main definitive host of *C. plica* in Denmark. The *C. plica* prevalence of Danish red foxes correspond with a study from Germany with a prevalence of 78% in red foxes (Bork-Mimm and Rinder, 2011). However, the prevalence in our study was lower than in Lithuania (98.3%) and higher than in Norway (53.0%) and Hungary (52.0%) (Bork-Mimm and Rinder, 2011; Bružinskaitė-Schmidhalter et al., 2012; Davidson et al., 2006; Magi et al., 2015; Saeed et al., 2006). In addition, red foxes are becoming increasingly more “urbanized” wildlife (Deplazes et al., 2004; Gloor et al., 2002; Scott et al., 2014), carrying the parasite close to premises where pet animals are roaming. Thus, red foxes are seemingly the most important factor in the epidemiology and risk of spread of the parasite to pet dogs.

In our study, the *C. plica* prevalence in raccoon dogs (0.5%) is considerably lower than in red foxes (73.7%). This observation correspond to a Lithuanian study documenting a prevalence of 11.3% in raccoon dogs and 93.3% in red foxes (Bružinskaitė-Schmidhalter et al., 2012). To our knowledge, the Lithuanian study is the only other study where *C. plica* infection in raccoon dogs have been analysed.

Ingestion of infected earthworms is the sole documented infection method for *C. plica* in canines. Elmeros et al. (2018) and Mikkelsen et al. (2016) demonstrated that Danish raccoon dogs to a large extent eat earthworms (68.6% and 34.7% of animals). In contrast, red foxes rarely eat earthworms, but prefer microtine rodents (Kauhala et al., 1998; Pagh et al., 2015a, 2015b). Hence, the *C. plica* prevalence herein observed for both raccoon dogs and red foxes contradicts the feed preference. This could indicate that raccoon dogs are less suitable as final host for *C. plica*. This applies for badgers, too. A Spanish study documented a prevalence of 2.1% in Eurasian badgers (Torres et al., 2001), although badgers frequently consume earthworms (Kauhala et al., 1998; Madsen et al., 2002). In addition to acquire the infection through consumption of earthworms, it is likely that red foxes can acquire the infection through paratenic hosts like rodents and birds. When rodents and birds consume *C. plica*-infected earthworms, the parasite seemingly enters these animals without undergoing any further development, but might remain alive till it gains entry to the red fox through consumption of the paratenic host. However, further studies are needed to identify paratenic hosts. A second possibility is that another, hitherto undocumented intermediate host exist. Lastly, *C. plica* infection might accumulate in red fox, so once infected, the parasite reside within the bladder for an extended time. This could be the case if an age effect was

**Table 1**  
*Capillaria plica* egg load in urine sediment or sediment from washing of the bladder.

Species	No examined	No positive animals (%)	95% confidence interval	1-10 eggs per ml (%)	11-20 eggs per ml (%)	21-30 eggs per ml (%)	> 30 eggs per ml (%)
Red foxes ( <i>Vulpes vulpes</i> )	247	182 (73.7)	68.2–79.2	41 (23.0)	31 (17.4)	12 (6.7)	94 (52.8)
Raccoon dogs ( <i>Nyctereutes procyonoides</i> )	375	2 (0.5)	0.2–1.3	1 (50.0)	1 (50.0)	–	–
Golden jackal ( <i>Canis aureus</i> )	1	1 (100.0)	–	–	–	–	1 (100.0)
Beech martens ( <i>Martes foina</i> )	20	4 (20.0)	0.8–39.2	–	4 (100.0)	–	–




Fig. 2. The prevalence of *Capillaria plica* infections in red foxes per region of Denmark (positive red foxes/total number of red foxes examined). The origin of five red foxes was unknown.

noted. Unfortunately, the age was not recorded in this study. A last possibility is that *Capillaria* eggs documented from the various hosts are indeed different species. However, our study was limited by lack of molecular analysis of *Capillaria* eggs from the various species.

The red foxes from Jutland had significantly higher *C. plica* prevalence compared to red foxes from Zealand. Jutland is the peninsula that forms the continental portion of Denmark. The population density is 84 persons per km<sup>2</sup>. In comparison, the population density is 358 persons per km<sup>2</sup> on Zealand. Since only few red foxes originated from Zealand, we refrain to conclude that the difference in prevalence between regions is affected by the availability of infected earthworms or the feeding habits of red foxes.

This is the first report of *C. plica* infection in beech martens (4 positive out of 20). *Capillaria plica* have been recorded in American martens with a prevalence of 6% (Seville and Addison, 1995), which determine that the genus *martes* are final hosts of *C. plica*.

In summary, our findings confirm that most Danish red foxes excrete *C. plica* eggs and that red foxes are the most likely reservoir host for infection in dogs. Moreover, further studies are needed to identify other possible intermediate hosts and infection through transport hosts.

#### Acknowledgements

This work was supported by the Ministry of Environment and Food of Denmark [grant numbers MST-854-00060]. A special thanks to laboratory technicians Boi Tien Thi Pharm and Aleksandra Tofteby. Without their hard work, this study was not a reality.

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ijppaw.2018.09.006>.

[doi.org/10.1016/j.ijppaw.2018.09.006](https://doi.org/10.1016/j.ijppaw.2018.09.006).

#### References

- Bork-Mimm, S., Rinder, H., 2011. High prevalence of *Capillaria plica* infections in red foxes (*Vulpes vulpes*) in Southern Germany. *Parasitol. Res.* 108, 1063–1067. <https://doi.org/10.1007/s00436-010-2196-0>.
- Bružinskaitė-Schmidhalter, R., Šarkūnas, M., Malakauskas, A., Mathis, A., Torgerson, P.R., Deplazes, P., 2012. Helminths of red foxes (*Vulpes vulpes*) and raccoon dogs (*Nyctereutes procyonoides*) in Lithuania. *Parasitology* 139, 120–127. <https://doi.org/10.1017/S0031182011001715>.
- Butterworth, E.W., Beverley-Burton, M., 1980. The taxonomy of *Capillaria* spp. (Nematoda: Trichuroidea) in carnivorous mammals from Ontario, Canada. *Syst. Parasitol.* 1, 211–236. <https://doi.org/10.1007/BF00009847>.
- Callegari, D., Kramer, L., Cantoni, A.M., Di Lecce, R., Dodi, P.L., Grandi, G., 2010. Canine bladderworm (*Capillaria plica*) infection associated with glomerular amyloidosis. *Vet. Parasitol.* 168, 338–341. <https://doi.org/10.1016/J.VETPAR.2009.11.008>.
- Davidson, R.K., Gjerde, B., Vikøren, T., Lillehaug, A., Handeland, K., 2006. Prevalence of *Trichinella* larvae and extra-intestinal nematodes in Norwegian red foxes (*Vulpes vulpes*). *Vet. Parasitol.* 136, 307–316. <https://doi.org/10.1016/J.VETPAR.2005.11.015>.
- Deplazes, P., Hegglin, D., Gloor, S., Romig, T., 2004. Wilderness in the city: the urbanization of *Echinococcus multilocularis*. *Trends Parasitol.* 20, 77–84.
- Elmeros, M., Malene, D., Mikkelsen, G., Nørgaard, L.S., Pertoldi, C., Hammer Jensen, T., Chriél, M., 2018. The diet of feral raccoon dog (*Nyctereutes procyonoides*) and native badger (*Meles meles*) and red fox (*Vulpes vulpes*) in Denmark. *Mamm. Res.* 1–9. <https://doi.org/10.1007/s13364-018-0372-2>.
- Gloor, S., Von, B., Ziswiler, V., Breitenmoser, U., Funk, S., 2002. The rise of urban foxes (*Vulpes vulpes*) in Switzerland and ecological and parasitological aspects of a fox population in the recently colonised city of Zurich. *Zeitschrift für Säugetierkd.* 66, 155–164.
- Kauhala, K., Laukkanen, P., von Rége, I., 1998. Summer food composition and food niche overlap of the raccoon dog, red fox and badger in Finland. *Ecography* 21, 457–463. <https://doi.org/10.1111/j.1600-0587.1998.tb00436.x>.
- Lamm, J., Main, F.A., 1964. Der Parasitenbefall bei Rotfischchen in Südhessen. *Z. Jagdwiss.* 10, 137–142.
- Magi, M., Guardone, L., Prati, M.C., Mignone, W., Macchioni, F., 2015. Extraintestinal nematodes of the red fox *Vulpes vulpes* in north-west Italy. *J. Helminthol.* 89, 506–511. <https://doi.org/10.1017/S0022149X1400025X>.
- Madsen, S.A., Madsen, A.B., Elmeros, M., 2002. Seasonal food of badgers (*Meles meles*) in

- Denmark. Mammalia 3, 341–352. <https://doi.org/10.1515/mamm.2002.66.3.341>.
- Mikkelsen, D.M.G., Nørgaard, L.S., Jensen, T.H., Chriél, M., Pertoldi, C., Elmeros, M., 2016. Mårhundens (*Nyctereutes procyonoides*) føde og fødeoverlap med hjemmehørende rovdyr i Danmark [In Danish]. Flora og Fauna 122, 101–114.
- Pagh, S., Skjold Tjørnløv, R., Illemann, J.K., Tolsgaard, S., Chriel, M., 2015a. Habitatrelateret føde hos ræv (*Vulpes vulpes*) i land-brugsområder. Flora Fauna 121, 1–14.
- Pagh, S., Skjold Tjørnløv, R., Olesen, C.R., Chriel, M., 2015b. The diet of Danish red foxes (*Vulpes vulpes*) in relation to a changing agricultural ecosystem. A historical perspective. Mammal Res 60, 319–329. <https://doi.org/10.1007/s13364-015-0244-y>.
- Saeed, I., Maddox-Hyttel, C., Monrad, J., Kapel, C.M.O., 2006. Helminths of red foxes (*Vulpes vulpes*) in Denmark. Vet. Parasitol. 139, 168–179. <https://doi.org/10.1016/j.vetpar.2006.02.015>.
- Scott, D.M., Berg, M.J., Tolhurst, B.A., Chauvenet, A.L.M., Smith, G.C., Neaves, K., Lochhead, J., Baker, P.J., 2014. Changes in the distribution of red foxes (*Vulpes vulpes*) in urban areas in Great Britain: findings and limitations of a media-driven nationwide survey. PLoS One 9, e99059. <https://doi.org/10.1371/journal.pone.0099059>.
- Senior, D.F., Solomon, G.B., Goldschmidt, M.H., Joyce, T., Bovee, K.C., 1980. *Capillaria plica* infection in dogs. J. Am. Vet. Med. Assoc. 176, 901–905.
- Seville, R.S., Addison, E.M., 1995. Nongastrointestinal helminths in marten (*Martes americana*) from Ontario, Canada. J. Wildl. Dis. 31, 529–533. <https://doi.org/10.7589/0090-3558-31.4.529>.
- Sréter, T., Széll, Z., Marucci, G., Pozio, E., Varga, I., 2003. Extraintestinal nematode infections of red foxes (*Vulpes vulpes*) in Hungary. Vet. Parasitol. 115, 329–334. [https://doi.org/10.1016/S0304-4017\(03\)00217-6](https://doi.org/10.1016/S0304-4017(03)00217-6).
- Torres, J., Miquel, J., Motjea, M., 2001. Helminth parasites of the eurasian badger (*Meles meles* L.) in Spain: a biogeographic approach. Parasitol. Res. 87, 259–263. <https://doi.org/10.1007/s004360000316>.